



Using a patient prompt list to raise concerns in oncology clinics does not necessarily lead to longer consultations

Rogers, S., Semple, C., Humphris, G., Lowe, D., & Kanatas, A. (2020). Using a patient prompt list to raise concerns in oncology clinics does not necessarily lead to longer consultations. *British Journal of Oral and Maxillofacial Surgery*, 58(9), 1164-1171. <https://doi.org/10.1016/j.bjoms.2020.08.035>

[Link to publication record in Ulster University Research Portal](#)

Published in:

British Journal of Oral and Maxillofacial Surgery

Publication Status:

Published (in print/issue): 01/11/2020

DOI:

<https://doi.org/10.1016/j.bjoms.2020.08.035>

Document Version

Author Accepted version

General rights

Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.

Using a patient prompt list to raise concerns in oncology clinics does not necessarily lead to longer consultations

Simon N Rogers, Cherith Semple, Gerald M Humphris, Derek Lowe, Anastasios Kanatas

Professor Simon N Rogers Faculty of Health and Social Care, Edge Hill University, Ormskirk, L39 4QP and Liverpool Head and Neck Centre, Liverpool University Hospital Aintree, Liverpool, UK
simonn.rogers@aintree.nhs.uk ORCID: 0000-0002-5989-6142

Dr Cherith Semple Institute of Nursing and Health Research, Ulster University, Shore Road, Newtownabbey, Co. Antrim, BT37 0QB and South Eastern Health & Social Care Upper Newtownards Road, Belfast, BT16 1RH. cherith.semple@setrust.hscni.net
ORCID: 0000-0002-4560-7637

Professor Gerald M Humphris School of Medicine, Medical & Biological Sciences, North Haugh, St Andrews, UK gmh4@st-andrews.ac.uk

Mr Derek Lowe Medical Statistician, Director, Astraglobe Ltd, Congleton, Cheshire.
astraglobeltd@btconnect.com

Professor Anastasios Kanatas³ Leeds Teaching Hospitals and St James Institute of Oncology, Leeds Dental Institute and Leeds General Infirmary, Leeds, UK
anastasios.kanatas@nhs.net

Corresponding author

Professor Simon N Rogers FRCS FRCS (maxfac) MD, Liverpool Head and Neck Centre, Liverpool University Hospital Aintree, Liverpool, UK
simonn.rogers@aintree.nhs.uk / snrogers.aintree@gmail.com Tel : 0151 529 5287

Keywords; head and neck cancer; intervention; prompt list; health related quality of life; randomised trial; Patient Concerns Inventory; consultations; duration

Using a patient prompt list to raise concerns in oncology clinics does not necessarily lead to longer consultations

Abstract

Head and Neck oncology post-treatment consultations form a critical component of care in terms of support and surveillance. They occur frequently in the first few years and can place substantial demands on health care resources. However, they provide useful opportunities for patients to raise issues and receive tailored information and support. The aim of this paper is to assess whether the use of a 56 item patient prompt list (PCI), completed immediately prior to the consultation significantly increases its duration. This was a pragmatic cluster preference randomised control trial of 288 patients with 15 consultant clusters from two sites either 'using' (n=8) or 'not using' (n=7) the PCI. Consultation times were known for 283 patients (136 PCI, 147 non-PCI) who attended their first post-treatment trial consultation, a median (IQR) 103 (70-160) days after the end of treatment. Consultations lasted a median (IQR) of 10 (7-13) minutes, mean 11.1 in non-PCI patients and a median (IQR) of 11 (8-15) minutes, mean 12.0 in PCI patients (p=0.07). After adjustment for patient clustering and significant case-mix the 95% confidence interval for the mean difference was from 1.45 minutes shorter with PCI to 2.98 minutes longer, p=0.50. There was significant variation in duration by consultant, tumour stage, treatment mode, overall QOL, distress (all p<0.001). In those completing the PCI, duration increased with the total number of PCI items selected (p<0.001). In conclusion, the inclusion of a prompt list to help facilitate the conversation with patients did not make a substantial difference to consultation times.

Keywords: head and neck cancer; intervention; prompt list; health related quality of life; randomised trial; Patient Concerns Inventory; consultations; duration

Introduction

There are considerable demands on head and neck cancer (HNC) clinic capacity which are already under pressure to meet the two-week suspected cancer pathway, provide time to discuss the diagnosis and treatment options for new patients and to review patients after treatment. The burden is likely to grow due to increasing rates of HNC and more patients under follow-up for late effects. Although practices vary there is a higher frequency of follow-up in the first two years post treatment. In the first year this is between one to three months, and in the second year two to four months.¹ The schedule of review visits can be stratified by risk of loco-regional recurrence, with high risk patients being seen more frequently.^{2,3} However, the follow-up consultation is valuable for several reasons. These include the assessment of treatment response, the identification of recurrence or new primary tumours, the monitoring and management of complications, the checking for physical and psychosocial consequences, the opportunity to address unmet needs and provide support to patients and their families, and a chance to deliver life-style advice (the 'teachable moment').

3-5

There are many issues that patients might wish to raise for discussion at their post treatment consultations.⁶ Patients value the opportunity to raise concerns,⁷ however report barriers to expressing these.⁸ Clinicians behaviour is the most dominant barrier, notably patients not being explicitly invited by the clinician to express concerns, and this is made worse in a setting of perceived lack of time.⁸ It is feasible to suggest that a prompt list approach, such as the Patient Concerns Inventory (PCI) might help reduce barriers.^{9,10} The PCI is a 56-item prompt list¹¹ and following a systematic review of various tools, it was recommended in the HNC setting due to excellent content validity.¹² Specialist HNC nurses and allied health professional when surveyed reported that of various tools available to assist in follow-up,

holistic needs assessment and survivorship care, they felt a HNC specific tool such as the PCI was more appropriate.¹³

When considering the added benefit of using the PCI in routine review consultations, clinicians might conversely be apprehensive about the additional time it might take when patients are afforded the opportunity to raise numerous issues. The PCI could significantly add to the length of the consultation and introduce delay in an already busy outpatient clinic. Hence, the main aim of this paper is to assess whether the use of the PCI completed immediately prior to the consultation significantly increases its duration. A secondary aim is to document other associations with duration of consultation, with particular focus on the number and nature of the items selected by patients using the PCI.

Methods

The data come from a pragmatic cluster-controlled trial conducted at two UK Cancer Centres, namely Aintree and Leeds. Consultants (the clustering factor) were randomised to ‘using’ or ‘not using’ an intervention incorporating the PCI prompt list at all their trial clinics. The methods have been described previously.¹⁴ Eligible patients were treated curatively for primary or secondary HNC, and all sites, stage of disease and treatments were included. Palliation and recurrence were exclusion criteria as were cognitive impairment, psychoses or dementia. The PCI consists of 56 clinical items¹¹ which patients select from before their appointment, to help guide the outpatient consultation through the symptoms and problems experienced following treatment for HNC. Patients were first discussed at MDT meetings between January 2017 and December 2018, with clinics between April 2017 and October 2019. The first post-treatment trial consultation with consultant surgeon is the focus of this paper.

Researchers at the Cancer Centres collected clinical and demographic data which matched those included in the head and neck 5000 project.¹⁸ HRQOL data were completed electronically (desktop, tablet, iPad) apart from one non-PCI consultant who used paper-based. HRQOL data included UW-QOLv4,¹⁵ Distress thermometer¹⁹ and EQ-5D-5L.²⁰ The UW-QOLv4 questionnaire consists of 12 single question domains, these having between 3 and 5 response options that are scaled evenly from 0 (worst) to 100 (best) according to the hierarchy of response.¹⁵ In regard to overall QOL, patients are asked to consider not only physical and mental health, but also many other factors, such as family, friends, spirituality or personal leisure activities that were important to their enjoyment of life. Subsequent analysis has led to the development of subscale composite scores¹⁶ and domain algorithms to screen for significant problems/dysfunction.¹⁷

Numerical data were summarised using the mean, median and Inter-quartile range (IQR). Boxplots showed variation in consultation times with boxplot and bar width scaled according to sample size. Asterisks represent values >3 box lengths from the edge of the box, with circles 1.5-3 box lengths away. Consultation times were compared using the Mann-Whitney (2 groups) or Kruskal-Wallis test (>2 groups). Spearman's correlation coefficient (r_s) was used to assess association between consultation time and other numerical/ordinal variables. Linear regression (STATA regress procedure) was used to estimate the unadjusted difference between PCI and non-PCI group mean consultation times and random effects linear regression (STATA xtreg procedure) was used to adjust for case-mix variables as independent predictors and for consultant clustering. To address skewness in the consultation times, standard errors (SEs) were estimated by a cluster bootstrap (5000 replications) that resampled with replacement over consultant clusters (rather than over individual

observations). Statistical significance was regarded as $p < 0.05$. SPSS v25 and Stata v13 were used for the analyses.

The PCI trial has ethical approval from North West - Liverpool Central Research Ethics Committee REC reference: IRAS 16/NW/0465, Project ID: 189554. It also has approval from the Health Research Authority (HRA). The Research and Development Department at [REDACTED] is coordinating the trial and [REDACTED] is the sponsor for the trial.

Results

Consultation times were known for 283 (Aintree 175, Leeds 108) of 288 attending their first post-treatment clinic, median (IQR) 193 (122-245) days after diagnosis and 103 (71-160) days after end of treatment. Fifteen consultants participated, seeing a median (IQR) of 16 (13-26) patients, range 5-48. Median (IQR) patient age was 62 (55-69) years and 69% (195) were male. Other patient characteristics are shown in Table 1.

Consultations lasted a median (IQR) 10 (8-14) minutes, mean 11.5, range 2 to 41. Aintree consultations took a median (IQR) of 10 (7-12) minutes, mean 10.1, compared with 12 (9-17), 13.9 at Leeds ($p < 0.001$). For PCI group patients they took a median (IQR) of 11 (8-15) minutes, mean 12.0, compared with 10 (7-13), 11.1 for Non-PCI patients ($p = 0.07$). The median time was longer for PCI patients by 1 minute at Aintree and 2 minutes at Leeds (Figure 1), with means of 10.5 (PCI) and 9.7 (Non-PCI) minutes at Aintree and 14.0 (PCI) and 13.9 (Non-PCI) minutes at Leeds. Individual consultants varied considerably (Figure 2), with means ranging from 7.0 to 16.4 minutes, medians from 7 to 15 minutes ($p < 0.001$).

Patients with advanced tumours, patients with treatments involving radiotherapy and/or chemotherapy and surgical patients having free-flap transfer had longer consultations (Table 1), as did patients living with others rather than living alone. Similar findings were seen for each centre (results not shown). There was no evidence of association between consultation time and days from the end of treatment to the clinic, $r_s=0.04$, $p=0.46$. Consultation time was weakly associated with the six-point overall QOL scale ($r_s=0.23$, $p<0.001$, Figure 3) and with the distress thermometer 0-10 scale ($r_s=0.21$, $p<0.001$, Figure 4), indicating longer consultation times with worse overall QOL and with greater distress. There were weak associations also with the UW-QOL physical function ($r_s=-0.21$, $p<0.001$) and social-emotional ($r_s=-0.23$, $p<0.001$) subscale scores and with the EQ-5D visual analog scale ($r_s=-0.16$, $p=0.008$), these indicating longer consultations with worse scale scores. Similar trends were noted for other HRQOL measures, with slightly longer consultation times for those registering UW-QOL domain dysfunction and shorter times for those giving the best UW-QOL domain response, as well as longer times for patients having moderate or more severe problems on each of the EQ-5D domains (results not shown).

In PCI group patients the median (IQR) number of PCI items selected was 5 (2-9), range 0-28, $n=136$. The top ten issues patients selected were dry mouth (50%, 68), dental health (35%, 48), chewing/eating (34%, 46), fear of recurrence (34%, 46), salivation (34%, 46), fatigue/tiredness (29%, 39), swallowing (29%, 39), taste (27%, 37), sore mouth (24%, 33) and mucus (24%, 33). Only two items were not selected: 'regret about treatment' and sexuality. Length of consultation increased with the total number of PCI items selected ($r_s=0.35$, $p<0.001$), this being driven by the numbers of items selected in the physical & functional, treatment-related and psychological / emotional / spiritual domains (Table 2). For

44 of the 56 PCI items both mean and median times were increased if that item was selected. Some of these increases were statistically significant (Table 2).

Linear regression (STATA regress) estimated the unadjusted difference between PCI and non-PCI group mean consultation times as 0.85 minutes longer in the PCI group (95% CI: 2.25 longer to 0.55 shorter), $p=0.23$. After adjustment for consultant clustering using random effects linear regression (STATA xtreg) and for case-mix factors of treatment, overall clinical stage, hospital site and living status (categories as per Table 1) as independent predictors, this became 0.54 minutes longer (95% CI: 2.75 longer to 1.66 minutes shorter), $p=0.63$. After further adjustment for overall QOL, distress thermometer, and physical and social-emotional composite scores (as per Table 1) it was 0.76 minutes longer (95% CI: 2.98 longer to 1.45 minutes shorter), $p=0.50$. The intraclass correlation (ICC) value estimate was 0.13.

Discussion

This paper spotlights the length of individual HNC patient clinic review consultations. It is appropriate to consider the frequency of follow-up consultations ² and also how to make this more time efficient without losing overall efficacy. This study is the first to detail duration of review consultations and the impact of including the PCI. The study has strengths, notably having 15 different consultants across two centres and a comparison of PCI use using the rigour of a RCT. However, the study is limited by the absence of other factors that could alter consultation length, such as consultant style and personality, number of carers in the consultation, timing of the consultation within the clinic schedule (beginning or end), and late running of consultations. Also, no record was made of the discussions that took place and how long it took to discuss specific issues. The PCI results need to be interpreted with caution data because items identified more commonly are more likely to reach statistical

significance and these are not necessarily those with the biggest observed time differences.

The study focuses on one clinic window around three months after treatment and as yet there is no data on how consultation lengths might vary over time. The study only included consultants and the length of consultations for surgeons in training was not assessed. The eight PCI consultants received one training session on using the PCI, either as a group presentation and discussion taking an hour or by individual training taking about 20 minutes. Anecdotally they reported how straightforward it was to include it as part of their usual practice but no attempt was made to evaluate this.

In this study, the addition of the PCI only led to a small increase in consultation length. This finding is supported by observation of clinicians in primary care, with patients having the opportunity to raise major concerns during the consultation, with no effect on visit length.²¹ The PCI enables patients to raise concerns that might otherwise be missed^{9, 10} and this can result in better patient satisfaction, improved communication and psychological wellbeing. In this current study, most PCI issues related to physical function, such as dry mouth, dental related concerns, and fatigue/tiredness, these forming a major component of clinic conversations. A common psychological issue is fear of recurrence. It was not known how long was spent addressing each issue; nevertheless, it is important to recognise the content as this helps to shape service delivery and policy making, as often the reality is that patients can wait a long time to access restorative dentistry and emotional support services.²² Perhaps, with further patient and clinician training and additional multi-professional support material, consultation times could be reduced without diminishing the therapeutic benefit. However, consultations are actually quite short and rather than focus solely on time efficiency the focus should perhaps be on reconfiguring services to address unmet need, enhancing HRQOL and the ‘teachable moment’ of doctor-patient interaction.

Advance tumour stage was associated with longer consultations, not surprising as these patients have more aspects of dysfunction to assess and discuss. Also, stage is associated with worse overall QOL, lower mood, and greater distress and these are associated with consultation duration. Furthermore, the total number of PCI items raised is associated with stage of cancer,⁶ low mood and poorer HRQOL.²³ The data shows considerable variability by individual consultant and this probably reflects distinct patterns of dialogue used in the doctor-patient communication. Given the complexity of HNC and its aftermath, and the reassurance of the physical examination, it is notable that most consultations were just over 10 minutes. This probably reflects time constraints inherent in busy out-patient clinics and the need to undertake reviews as time efficiently as possible. It is recognised that some consultations might need to be much longer to perform adequate review and address patient concerns. In light of this, there is an emergence and recognition of the extended role of CNSs and within many HNC outpatient clinics, follow-up care is supported or shared to promote holistic, personalised patient care.

In conclusion, the inclusion of a prompt list to help facilitate the conversation with patients does not seem to make a substantial difference to consultation times. As most patients really appreciate this holistic approach to their care, worries about its inclusion into routine consultations in respect to disrupting the flow of clinics is unfounded.

Funding sources

This trial is funded by the [REDACTED]

Conflict of Interest

No conflict of interest

Ethics statement/confirmation of patient permission

Ethical approval has been given. Patient permission not relevant

Acknowledgements

This paper presents independent research funded by the [REDACTED]

[REDACTED]

[REDACTED] The views expressed are those of the authors and not necessarily those of the [REDACTED]

References

- 1.Digonnet A, Hamoir M, Andry G, Vander Poorten V, Haigentz M Jr, Langendijk JA, de Bree R, Hinni ML, Mendenhall WM, Paleri V, Rinaldo A, Werner JA, Takes RP, Ferlito A. Follow-up strategies in head and neck cancer other than upper aerodigestive tract squamous cell carcinoma. *Eur Arch Otorhinolaryngol*. 2013 Jul;270(7):1981-9. doi: 10.1007/s00405-012-2220-3. Epub 2012 Oct 26.
- 2.Kanatas A, Bala N, Lowe D, Rogers SN. Outpatient follow-up appointments for patients having curative treatment for cancer of the head and neck: are the current arrangements in need of change? *Br J Oral Maxillofac Surg*. 2014 Oct;52(8):681-7. doi: 10.1016/j.bjoms.2014.06.017. Epub 2014 Jul 15.
- 3.Simo R, Homer J, Clarke P, Mackenzie K, Paleri V, Pracy P, Roland N. Follow-up after treatment for head and neck cancer: United Kingdom National Multidisciplinary Guidelines. *J Laryngol Otol*. 2016 May;130(S2):S208-S211.
- 4.Simcock R, Simo R. Follow-up and Survivorship in Head and Neck Cancer. *Clin Oncol (R Coll Radiol)*. 2016 Jul;28(7):451-8. doi: 10.1016/j.clon.2016.03.004. Epub 2016 Apr 16.

5. Nguyen NA, Ringash J. Head and Neck Cancer Survivorship Care: A Review of the Current Guidelines and Remaining Unmet Needs. *Curr Treat Options Oncol*. 2018 Jul 9;19(8):44. doi: 10.1007/s11864-018-0554-9.
6. Kanatas A, Ghazali N, Lowe D, Udberg M, Heseltine J, O'Mahony E, Rogers SN. Issues patients would like to discuss at their review consultation: variation by early and late stage oral, oropharyngeal and laryngeal subsites. *Eur Arch Otorhinolaryngol*. 2013 Mar;270(3):1067-74. doi: 10.1007/s00405-012-2092-6. Epub 2012 Jun 29.
7. Rogers SN, Lowe D. An evaluation of the Head and Neck Cancer Patient Concerns Inventory across the Merseyside and Cheshire Network. *Br J Oral Maxillofac Surg*. 2014 Sep;52(7):615-23. doi: 10.1016/j.bjoms.2014.04.011. Epub 2014 Jun 11
8. Brandes K, Linn AJ, Smit EG, van Weert JC. Patients' reports of barriers to expressing concerns during cancer consultations. *Patient Educ Couns*. 2015 Mar;98(3):317-22. doi: 10.1016/j.pec.2014.11.021. Epub 2014 Dec 2.
9. Ghazali N, Roe B, Lowe D, Rogers SN. Uncovering patients' concerns in routine head and neck oncology follow up clinics: an exploratory study. *Br J Oral Maxillofac Surg*. 2013 Jun;51(4):294-300. doi: 10.1016/j.bjoms.2012.08.002. Epub 2012 Aug 24.
10. Ghazali N, Roe B, Lowe D, Rogers SN. Patients concerns inventory highlights perceived needs and concerns in head and neck cancer survivors and its impact on health-related quality of life. *Br J Oral Maxillofac Surg*. 2015 Apr;53(4):371-9. doi: 10.1016/j.bjoms.2015.01.022. Epub 2015 Feb 20.
11. Rogers SN, El-Sheikha J, Lowe D. The development of a Patients Concerns Inventory (PCI) to help reveal patients concerns in the head and neck clinic. *Oral Oncol*. 2009 Jul;45(7):555-61. doi: 10.1016/j.oraloncology.2008.09.004. Epub 2008 Nov 22.
12. Shunmugasundaram C, Rutherford C, Butow PN, Sundaresan P, Dhillon HM. Content comparison of unmet needs self-report measures used in patients with head and neck cancer:

A systematic review. *Psychooncology*. 2019 Dec;28(12):2295-2306. doi: 10.1002/pon.5257.

Epub 2019 Nov 6.

13. Wells M, Semple CJ, Lane C. A national survey of healthcare professionals' views on models of follow-up, holistic needs assessment and survivorship care for patients with head and neck cancer. *Eur J Cancer Care (Engl)*. 2015 Nov;24(6):873-83. doi: 10.1111/ecc.12285.

Epub 2015 Jan 23.

14. Rogers SN, Lowe D, Lowies C, Yeo ST, Allmark C, Mcavery D, Humphris GM, Flavel R, Semple C, Thomas SJ, Kanatas A. Improving quality of life through the routine use of the patient concerns inventory for head and neck cancer patients: a cluster preference randomized controlled trial. *BMC Cancer*. 2018 Apr 18;18(1):444. doi: 10.1186/s12885-018-4355-0.

15. Rogers SN, Gwanne S, Lowe D, Humphris G, Yueh B, Weymuller EA Jr. The addition of mood and anxiety domains to the University of Washington quality of life scale. *Head Neck* 2002;24(6):521-9.

16. Rogers SN, Lowe D, Yueh B, Weymuller EA Jr. The physical function and social-emotional function subscales of the University of Washington Quality of Life Questionnaire. *Arch Otolaryngol Head Neck Surg* 2010;136:352-7. doi: 10.1001/archoto.2010.32.

17. Rogers SN, Lowe D. Screening for dysfunction to promote multidisciplinary intervention by using the University of Washington Quality of Life Questionnaire. *Arch Otolaryngol Head Neck Surg* 2009;135:369-75. doi: 10.1001/archoto.2009.7.

18. Ness AR, Waylen A, Hurley K, Jeffreys M, Penfold C, Pring M, Leary SD, Allmark C, Toms S, Ring S, Peters TJ, Hollingworth W, Worthington H, Nutting C, Fisher S, Rogers SN, Thomas SJ; Head and Neck 5000 Study Team. Recruitment, response rates and characteristics of 5511 people enrolled in a prospective clinical cohort study: head and neck 5000. *Clin Otolaryngol*. 2016 Dec;41(6):804-809. doi: 10.1111/coa.12548. Epub 2016 Feb 8.

- 19.Hegel MT, Collins ED, Kearing S, Gillock KL, Moore CP, Ahles TA. Sensitivity and specificity of the Distress Thermometer for depression in newly diagnosed breast cancer patients. *Psychooncology* 2008;17:556-60.
- 20.<https://euroqol.org/> [accessed June 2019].
- 21.Heritage J, Robinson JD, Elliott MN, Beckett M, Wilkes M. Reducing patients' unmet concerns in primary care: the difference one word can make. *J Gen Intern Med.* 2007 Oct;22(10):1429-33. Epub 2007 Aug 3.
- 22.Semple CJ, Lannon D, Qudairat E, McCaughan E, McCormac R. Development and evaluation of a holistic surgical head and neck cancer post-treatment follow-up clinic using touchscreen technology-Feasibility study.*Eur J Cancer Care (Engl).* 2018 Mar;27(2):e12809. doi: 10.1111/ecc.12809. Epub 2018 Feb 8.
- 23.Rogers SN, Alvear A, Anesi A, Babin E, Balik A, Batstone M, Brockmeyer P, Carrasco CC, Chien CY, Chirila M, Dholam KP, Doss JG, Finizia C, Ghani WMN, Gurav SV, Kadir K, Kolator M, Lima R, Lin YT, Nhongo S, Ozdemir-Karatas M, Peker K, Pesic Z, Ransy P, Santos I, Schliephake H, Shah K, Souza F, Sunil G, Thankappan K, Ehrsson YT, Tiple C, Tuomi L, Valerini S, Lara PV, Zatoński T, Lowe D. Variations in concerns reported on the patient concerns inventory in patients with head and neck cancer from different health settings across the world. *Head Neck.* 2020 Mar;42(3):498-512. doi: 10.1002/hed.26027. Epub 2019 Dec 13.

Table 1. Consultation times by case-mix factors

		Patients	Mean	Median (IQR)	P value*
	All patients	283	11.5	10 (8-14)	
Site	Aintree	175	10.1	10 (7-12)	<0.001
	Leeds	108	13.9	12 (9-17)	
Trial group	PCI	136	12.0	11 (8-15)	0.07
	Non-PCI	147	11.1	10 (7-13)	
Consultant experience	Before 2010	138	11.0	10 (8-12)	0.17
	Since 2010	145	12.0	11 (8-15)	
Gender	Female	88	11.6	10 (7-15)	0.72
	Male	195	11.5	10 (8-14)	
Age at clinic	<55	68	12.1	11 (9-15)	0.10
	55-64	115	11.0	10 (7-14)	
	65-74	67	12.5	11 (8-15)	
	≥75	33	10.3	9 (7-13)	
Tumour site:	Oral cavity	132	11.7	10 (7-15)	0.20
	Oropharynx	91	11.3	10 (8-14)	
	Larynx	38	10.9	9 (7-12)	
	Other	22	12.9	12 (9-16)	
Overall stage	Early 0-2	122	10.5	9 (7-12)	<0.001
	Advanced 3-4	161	12.3	11 (9-15)	
Primary treatment:	S only	93	9.3	9 (7-11)	<0.001
Surgery (S)	S only & FF	21	12.5	11 (7-15)	
RadioTherapy (RT)	RT or RT/CT only	57	12.1	10 (8-15)	
ChemoTherapy (CT)	S & (RT or RT/CT)	67	12.4	11 (9-15)	
Free Flap transfer (FF)	S & (RT or RT/CT) & FF	45	13.8	12 (8-19)	
WHO comorbidity	0	177	11.3	10 (8-13)	0.65
	1	65	12.3	10 (8-16)	
	2-4	41	11.5	10 (7-15)	
ACE27 comorbidity	None	135	11.4	10 (8-13)	0.60
	Mild	94	11.9	11 (8-15)	
	Moderate	47	11.1	10 (8-13)	
	Severe	7	13.6	14 (n/a)	
Living situation	Alone	65	9.9	9 (7-12)	0.01
In house/flat	With others	215	12.1	10 (8-15)	
Working	Yes	86	11.9	10 (8-14)	0.45
	No	189	11.5	10 (7-15)	
Financial benefits	Yes	105	11.6	10 (8-14)	0.44
	No	155	11.8	10 (8-15)	
Smoking habit	Current	37	10.8	9 (7-14)	0.15
	Former	158	11.6	10 (8-14)	
	Never	80	12.1	11 (8-15)	
Alcohol habit	Current	191	11.3	10 (7-14)	0.09
	Former	72	12.6	12 (8-15)	
	Never	13	10.6	10 (6-16)	
IMD 2019 quintile	1=least deprived	34	12.9	12 (9-15)	0.14
	2	55	12.2	11 (8-15)	
	3	48	11.3	10 (7-14)	
	4	39	11.9	11 (8-15)	
	5=most deprived	107	10.8	10 (7-12)	
Overall QOL**	Good, V good, Outstanding	195	10.8	10 (7-13)	0.004
	Fair, poor, V poor	88	13.2	11 (9-15)	
Social-emotional function score**	<60	71	13.5	11 (9-17)	0.01
	60-79	96	11.5	10 (8-14)	
	80-100	116	10.3	10 (7-13)	
Physical function Score**	<60	89	13.1	11 (8-17)	0.001
	60-79	107	11.5	11 (8-15)	
	80-100	87	10.0	9 (7-12)	
Distress Thermometer	<4	157	10.9	10 (7-13)	0.01
	≥4	126	12.3	11 (8-15)	

*Mann-Whitney test (2 comparison groups), Kruskal-Wallis (>2 comparison groups)

** From the UW-QOL v4

Table 2. Consultation times by the number of PCI items selected and by specific PCI items selected


	All patients	Patients 136	Mean 12.0	Median (IQR) 11 (8-15)	P value*
<u>Number of items selected</u>					
Total number	0-4	54	10.2	10 (7-12)	0.004
	5-9	49	12.5	11 (9-15)	
	10-14	20	13.8	15 (10-16)	
	≥15	13	14.4	15 (8-18)	
Physical & functional Wellbeing domain	0-4	72	10.2	10 (7-12)	<0.001
	5-9	42	14.1	14 (9-16)	
	≥10	22	13.7	12 (10-18)	
Treatment related domain	None	95	11.4	10 (8-14)	0.06
	≥1	41	13.2	12 (9-17)	
Social care and social Wellbeing domain	None	105	12.0	11 (8-15)	0.60
	≥1	31	12.0	11 (9-16)	
Psychological, emotional & spiritual domain	None	64	10.6	10 (7-13)	0.02
	One	42	12.6	12 (9-15)	
	≥2	30	14.1	12 (9-18)	
<u>Specific items selected (p<0.05)**</u>					
Depression	Yes	8	17.8	17 (n/a)	0.006
	No	128	11.6	11 (8-15)	
Mobility	Yes	10	17.1	15 (11-19)	0.01
	No	126	11.6	11 (8-15)	
Vomiting	Yes	5	16.6	18 (n/a)	0.01
	No	131	11.8	11 (8-15)	
Fatigue/tiredness	Yes	39	15.0	14 (11-18)	<0.001
	No	97	10.8	10 (8-14)	
Mucus	Yes	33	14.4	14 (9-19)	0.006
	No	103	11.2	10 (8-14)	
Dental health	Yes	48	13.8	14 (9-18)	0.01
	No	88	11.0	10 (8-13)	
Shoulder	Yes	30	14.0	12 (9-17)	0.03
	No	106	11.4	10 (8-14)	
Pain in head/neck	Yes	29	13.7	14 (10-16)	0.01
	No	107	11.5	10 (8-14)	
Fear of cancer returning	Yes	46	13.4	12 (9-16)	0.05
	No	90	11.3	10 (8-15)	
Salivation	Yes	46	13.1	12 (9-15)	0.01
	No	90	11.4	10 (8-14)	

*Mann-Whitney test (2 comparison groups), Kruskal-Wallis (>2 comparison groups)

** ordered according to the difference in mean times of when item selected and when not selected


n/a: IQR only computed if n≥10

Figure 1. Consultation times for PCI and non-PCI patients at Aintree and Leeds



Note that the boxplot and bar width are scaled according to the number of patients. An asterisk represents a value more than 3 box lengths from the upper or lower edge of the box, while a circle marks a value between 1.5 and 3 box lengths away from the box.


Figure 2. Consultation times by consultant



Aintree PCI: consultants 1, 2, 4, 6, 7
Aintree Non-PCI: consultants 3, 5, 8, 9
Leeds PCI: consultants 12, 15, 16
Leeds Non-PCI: consultants 11, 13, 14


Note that the boxplot and bar width are scaled according to the number of patients. An asterisk represents a value more than 3 box lengths from the upper or lower edge of the box, while a circle marks a value between 1.5 and 3 box lengths away from the box.

Figure 3. Consultation times by overall QOL stated by patients on the UW-QOLv4 questionnaire



Note that the boxplot and bar width are scaled according to the number of patients. An asterisk represents a value more than 3 box lengths from the upper or lower edge of the box, while a circle marks a value between 1.5 and 3 box lengths away from the box.

Figure 4. Consultation times by the Distress thermometer scale



Note that the boxplot and bar width are scaled according to the number of patients. An asterisk represents a value more than 3 box lengths from the upper or lower edge of the box, while a circle marks a value between 1.5 and 3 box lengths away from the box.